

resultant piRNAs. If we can monitor the behavior of each piRNA in cells, we will be able to show whether this pathway actually exists or not. Towards this goal, it will be necessary to develop a system in which piRNAs can be traced individually in their amplification cycle.

SUPPLEMENTAL INFORMATION

Supplemental Information contains one table and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2015.07.001>.

ACKNOWLEDGEMENTS

We thank Shizuo George Kamita for critical reading of the manuscript. The authors declare no competing financial interests.

REFERENCES

- Iwasaki, Y.W., Siomi, M.C., and Siomi, H. (2015). PIWI-interacting RNA: its biogenesis and functions. *Annu. Rev. Biochem.* 84, 405–433.
- Itou, D., Shiromoto, Y., Shin-ya, Y., Ishii, C., Nishimura, T., Ogonuki, N., Ogura, A., Hasuwa, H., Fujihara, Y., Kuramochi-Miyagawa, S., et al. (2015). Induction of DNA methylation by artificial piRNA production in male germ cells. *Curr. Biol.* 25, 901–906.
- Nishimasu, H., Ishizu, H., Saito, K., Fukuhara, S., Kamatani, M.K., Bonfond, L., Matsumoto, N., Nishizawa, T., Nakanaga, K., Aoki, J., et al. (2012). Structure and function of Zucchini endoribonuclease in piRNA biogenesis. *Nature* 491, 284–287.
- Ipsaro, J.J., Haase, A.D., Knott, S.R., Joshua-Tor, L., and Hannon, G.J. (2012). The structural biochemistry of Zucchini implicates it as a nuclease in piRNA biogenesis. *Nature* 491, 279–283.
- Kawaoka, S., Izumi, N., Katsuma, S., and Tomari, Y. (2011). 3' end formation of PIWI-interacting RNAs in vitro. *Mol. Cell.* 43, 1015–1022.
- Muerdter, F., Olovnikov, I., Molaro, A., Rozhkov, N.V., Czech, B., Gordon, A., Hannon, G.J., and Aravin, A.A. (2012). Production of artificial piRNAs in flies and mice. *RNA* 18, 42–52.
- Kawaoka, S., Mitsutake, H., Kikuchi, T., Kobayashi, M., Yohikawa, M., Suzuki, Y., Sugano, S., Shimada, T., Kobayashi, J., Tomari, Y., et al. (2012). A role for transcription from a piRNA cluster in de novo piRNA production. *RNA* 18, 265–273.
- Yamamoto, Y., Watanabe, T., Hoki, Y., Shirane, K., Li, Y., Ichiiyanagi, K., Kuramochi-Miyagawa, S., Toyoda, A., Fujiyama, A., Oginuma, M., et al. (2013). Targeted gene silencing in mouse germ cells by insertion of a homologous DNA into a piRNA generating locus. *Genome Res.* 23, 292–299.
- Hirano, T., and Siomi, H. (2015). Small RNAs: Artificial piRNAs for transcriptional silencing. *Curr. Biol.* 25, R280–R283.

Department of Agricultural and Environmental Biology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan.

*E-mail: katsuma@ss.ab.a.u-tokyo.ac.jp

Correspondence

Reply to Shoji and Katsuma

Satomi Kuramochi-Miyagawa¹
and Toru Nakano^{1,2,*}

Our paper reports a novel strategy for the artificial introduction of DNA methylation in mouse gonocytes [1]. The manuscript presents data showing that the concomitant expression of sense and anti-sense of EGFP transgenes in embryonic male germ cells induces gene silencing via the piRNA pathway and that the expression of an antisense Dnmt3L transgene induces silencing of the endogenous Dnmt3L gene.

Based on our study of EGFP transgenic mice, we concluded that the expression of sense and anti-sense EGFP genes is necessary and sufficient for piRNA production. Analyzing the data on piRNA expression from the Miwi2 promoter region of the antisense EGFP transgenic mice, Shoji and Katsuma raised a serious concern regarding this conclusion [2]. Given the data on the piRNAs of the Miwi2 promoter region, we would like to withdraw one of our conclusions, “concomitant expression of sense and antisense RNA transcripts is necessary and sufficient for piRNA production.”

We concluded based on gene capture analysis that the transgene was not integrated into a ‘typical’ piRNA cluster, because the number of piRNAs corresponding to the integrated region was not significant. That is, we adopted a kind of ‘prospective’ definition of piRNA clusters. Meanwhile, Shoji and Katsuma adopted criteria to define piRNA clusters based on bioinformatic analysis of transgenic mice. For transgenes that produce piRNAs corresponding to their own sequences, the loci into which the transgenes are integrated are interpreted as belonging to the piRNA clusters. We agree with their claim based on this definition; however, at the same time, we consider that a more precise definition of piRNA clusters would be necessary.

There are still some ambiguities to be addressed. Although we did not perform deep sequencing analysis of small RNAs, similar gene silencing by the antisense EGFP transgene was found in two other transgenic lines. These transgenic lines silenced the expression

of Oct4–EGFP in the double transgenic mice, in which both sense and antisense EGFP were expressed concomitantly, as well as transgenic line #1. In order to draw a robust conclusion, we will further analyze piRNA production in these two antisense transgenic animals and report the results in the near future.

There are two possible results. In the first case, these transgenic mice may also produce piRNAs corresponding to the MIWI2 promoter and EGFP transgene. If this is the case, we are willing to conclude that they are integrated into ‘minor’ or ‘atypical’ piRNA clusters. In this case, comparison of the expression of the amount of piRNA corresponding to the transgene will allow us to determine whether the amount of piRNAs or another factor is necessary for gene silencing via the piRNA pathway. This is because the simple existence of transgene-related piRNAs did not induce gene silencing in antisense transgenic mice (Figure 1C in [1]). In the second case, at least one of the two transgenic mouse lines may not produce a significant amount of piRNAs corresponding to the transgene. This would confirm our initial interpretation.

The main source of our misunderstanding was the discrepancy between artificial piRNA production and gene silencing. During the course of our experiments and publication process, we mistakenly believed that they should be concomitant. This discrepancy raises an interesting and important question about the relationship between piRNA production and gene silencing. Abundant piRNAs and/or piRNAs with some unknown characteristics may be necessary for DNA methylation and subsequent gene silencing. Finally, we would like to emphasize that all of the data presented in our paper and the main conclusion are robust.

REFERENCES

- Itou, D., Shiromoto, Y., Yukiho, S.Y., Ishii, C., Nishimura, T., Ogonuki, N., Ogura, A., Hasuwa, H., Fujihara, Y., Kuramochi-Miyagawa, S., and Nakano, T. (2015). Induction of DNA methylation by artificial piRNA production in male germ cells. *Curr. Biol.* 25, 901–906.
- Shoji, K., and Katsuma, S. (2015). Is the expression of sense and antisense transgenes really sufficient for artificial piRNA expression? *Curr. Biol.* 25, R708–R710.

Department of Pathology, ¹Medical School and ²Graduate School of Frontier Biosciences, Osaka University, Yamada-oka 2-2 Suita, Osaka 565-0871, Japan.

*E-mail: tnakano@patho.med.osaka-u.ac.jp